Linctagon®-C

Effervescents Tablets

SCHEDULING STATUS
To be allocated.

PROPRIETARY NAME AND DOSAGE FORM
Linctagon®-C (Effervescent Tablets)

COMPOSITION

Active Ingredients	Per effervescent tablet:	Per 3 effervescent tablets (max. daily dosage):	%RDA# per max daily dosage:
Pelargonium sidoides Root Extract	350 mg	1050 mg	*
Vitamin A	440 µg (800 IU)	1320 µg (2400 IU)	72
Vitamin C	500 mg	1500 mg	2500
MSM (Methylsulfonylmethane)	400 mg	1200 mg	*
Quercetin	80 mg	240 mg	*
Bromelain (80 GDU)	50 mg	150 mg	*
Zinc	5 mg	15 mg	100

#Recommended Daily Allowance for adults and children older than 10 years.
*RDA not established.
Contains artificial sweeteners (sodium saccharin and sodium cyclamate).

PHARMACOLOGICAL CLASSIFICATION
D.32.16 Other Substances or Agents - Others.
Combination Western Herbal Medicine.

INDICATIONS
Linctagon®-C contains the active ingredient *Pelargonium sidoides* that can assist the body with symptomatic relief associated with colds & flu. For a diagnosis or if symptoms persist, consult a medical practitioner. Efficacy of support may vary between users. Suitable for people with diabetes.

CONTRAINDICATION
Do not use if known hypersensitivity or allergy exists towards any of the ingredients. If in doubt consult your medical practitioner.

WARNINGS AND SPECIAL PRECAUTIONS
Do not use if known hypersensitivity or allergy exists towards any of the ingredients. If in doubt consult your medical practitioner. People using anticoagulant - or antiplatelet medication, such as Warfarin, should use this product with caution as *Pelargonium sidoides* may have a blood thinning effect. Products containing herbal ingredients may interact with some medication. If you are taking any chronic medication, do not use this product without consulting your medical practitioner. Do not exceed the recommended daily dosage. Do not use after date of expiry. Check expiry date on unit.
Porphyria: Safety has not been established.
KEEP OUT OF REACH OF CHILDREN.

INTERACTION
People using anticoagulant - or antiplatelet medication, such as Warfarin, should use this product with caution as *Pelargonium sidoides* may have a blood thinning effect.

PREGNANCY AND LACTATION
Safety for use during pregnancy and lactation (breastfeeding) has not been established.

DOSAGE AND DIRECTIONS FOR USE
Do not exceed the recommended daily dosage.
For optimal efficacy, take the recommended daily dosage for at least 5 days.
Adults and children 12 years and older:
Take one effervescent tablet, dissolved in half a glass of water, three times a day, after a meal.
For children under the age of 12, Linctagon®-C Junior is available in two flavours.

SIDE EFFECTS
If you experience any side effects or sensitivity towards any of the ingredients, discontinue use. If symptoms persist, or if any adverse reactions occur, consult a medical practitioner. No side effects have been reported.

KNOWN SYMPTOMS OF OVERDOSAGE AND PARTICULARS OF ITS TREATMENT
Do not exceed the recommended daily dosage. If an accidental overdose occurs, please consult a medical practitioner. Symptoms may include, but are not limited to, gastrointestinal disturbances and headaches. Treatment is symptomatic and supportive.

IDENTIFICATION
Different flavour variants available.
Orange flavour: Round, orange colour and orange flavour effervescent tablet.
Cherry flavour: Round, pink colour and cherry flavour effervescent tablet.

PRESENTATION
Plastic tube container, with 12 or 20 effervescent tablets, in a unit carton, with a package insert.

STORAGE INSTRUCTIONS
Store below 25 °C. Store away from moisture and direct sunlight.
Replace cap firmly after use.
KEEP OUT OF REACH OF CHILDREN.

REGISTRATION NUMBER
To be allocated.

NAME AND BUSINESS ADDRESS OF THE HOLDER OF THE CERTIFICATE OF REGISTRATION
Nativa (Pty) Ltd
260 Cradock Avenue, Lyttelton, Centurion, 0157,
Gauteng, South Africa
Tel: +27 (0) 12 664 7110 • Fax: +27 (0) 12 664 8031
Customer Care Line: **0860 628 482**
E-mail: health@nativa.co.za
Website: www.nativa.co.za

DATE OF PUBLICATION OF THE PACKAGE INSERT
May 2006

This unregistered medicine has not been evaluated by the SAHPRA for its quality, safety or intended use.

Linctagon® - C

Nativa health • science • nature

Bruistablette

SKEDULERING STATUS
Moet toegeken word.

EIENDOMSNAAM EN DOSEERVORM
Linctagon®-C (Bruistablette)

SAMESTELLING

Aktiewe Bestanddele	Per bruistablet:	Per 3 bruistablette (maks. daaglikse dosis):	%ADT# per maks. daaglikse dosis:
Pelargonium sidoides Wortelekstrak	350 mg	1050 mg	*
Vitamien A	440 µg (800 IU)	1320 µg (2400 IU)	72
Vitamien C	500 mg	1500 mg	2500
MSM (Metielsulfonielmetaan)	400 mg	1200 mg	*
Kwersitin	80 mg	240 mg	*
Bromelien (80 GDU)	50 mg	150 mg	*
Sink	5 mg	15 mg	100

#Aanbevole Daaglikse Toelaag vir volwassenes en kinders ouer as 10 jaar.
*ADT nie vasgestel nie.
Bevat kunsmatige versoeters (natriumsakkarien en natriumsiklamaat).

FARMAKOLOGIESE KLASSIFIKASIE
D.32.16 Ander Stowwe of Middels - Ander.
Kombinasie Westerse Kruimedisyne.

INDIKASIES
Linctagon®-C bevat die aktiewe bestanddeel *Pelargonium sidoides* wat die liggaam kan ondersteun met die verligting van simptome wat met verkoues & griep geassosieer word. Raadpleeg 'n geneesheer vir 'n diagnose of indien simptome voortduur. Effektiwiteit van ondersteuning kan verskil tussen gebruikers. Geskik vir persone met diabetes.

KONTRA-INDIKASIE
Persone met hipersensitiwiteit of allergieë vir enige van die bestanddele moet nie die produk gebruik nie. Indien twyfel bestaan, raadpleeg u geneesheer.

WAARSKUWINGS EN SPESIALE VOORSORGMAATREËLS
Persone met hipersensitiwiteit of allergieë vir enige van die bestanddele moet nie die produk gebruik nie. Indien twyfel bestaan, raadpleeg u geneesheer. Persone wat antikoagulant- of bloedverdunningsmedikasie, soos Warfarin neem, moet die produk met omsigtigheid gebruik, aangesien *Pelargonium sidoides* 'n bloedverdunningseffek kan hê. Produkte wat kruiebestanddele bevat, kan interaksies met sommige medikasie hê. Indien enige chroniese medikasie gebruik word, raadpleeg eers u geneesheer voordat die produk gebruik word. Moet nie die aanbevole daaglikse dosis oorskry nie. Die produk moet nie gebruik word nadat die vervaldatum reeds bereik is nie. Sien vervaldatum op die eenheid.
Porfirie: Veiligheid is nog nie vasgestel nie.
HOU BUITE BEREIK VAN KINDERS.

INTERAKSIE
Persone wat antikoagulant- of bloedverdunningsmedikasie, soos Warfarin neem, moet hierdie produk met omsigtigheid gebruik, aangesien *Pelargonium sidoides* 'n bloedverdunningseffek kan hê.

SWANGERSKAP EN LAKTASIE
Veiligheid vir gebruik tydens swangerskap en laktasie (borsvoeding) is nog nie vasgestel nie.

DOSIS EN GEBRUIKSAANWYSINGS
Moet nie die aanbevole daaglikse dosis oorskry nie.
Gebruik die aanbevole daaglikse dosis vir ten minste 5 dae vir optimale effektiwiteit.
Volwassenes en kinders 12 jaar en ouer:
Neem een bruistablet, opgelos in 'n halwe glas water, drie keer per dag, na 'n ete.
Linctagon®-C Junior is beskikbaar in twee geure vir kinders onder 12 jaar oud.

NEWE-EFFEKTE
Indien enige newe-effekte of sensitiwiteit teenoor enige van die bestanddele voorkom, staak gebruik. Indien simptome voortduur of nadelige reaksies ondervind word, raadpleeg 'n geneesheer.
Geen newe-effekte is gerapporteer nie.

BEKENDE SIMPTOME VAN OORDOSERING EN BEHANDELING
Moet nie die aanbevole daaglikse dosis oorskry nie. Raadpleeg 'n geneesheer indien 'n oordosis per abuis ingeneem word. Simptome kan insluit, maar is nie beperk tot hoofpyn en gastroïntestinale ongemak, nie. Behandeling is simptomaties en ondersteunend.

IDENTIFIKASIE
Verskillende geurvariante beskikbaar.
Lemoengeur: Ronde, oranje-kleurige, lemoengeur bruistablet.
Kersiegeur: Ronde, pienk-kleurige, kersiegeur bruistablet.

AANBIEDING
Plastiese buishouer met 12 of 20 bruistablette, in 'n eenheidskarton, met 'n voubiljet.

BERGINGSAANWYSINGS
Bewaar benede 25 °C. Bewaar weg van vog en direkte sonlig.
Plaas doppie stewig terug na gebruik.
HOU BUITE BEREIK VAN KINDERS.

REGISTRASIENOMMER
Moet toegeken word.

NAAM EN BESIGHEIDSADRES VAN DIE HOUER VAN DIE SERTIFIKAAT VAN REGISTRASIE
Nativa (Edms) Bpk
Cradocklaan 260, Lyttelton, Centurion, 0157, Gauteng, Suid-Afrika
Tel: +27 (0) 12 664 7110 • Faks: +27 (0) 12 664 8031
Verbruikershulplyn: **0860 628 482**
E-pos: health@nativa.co.za
Webblad: www.nativa.co.za

DATUM VAN PUBLIKASIE VAN DIE VOUBILJET
Mei 2006

Hierdie medisyne is nog nie deur die SAHPRA geregistreer of geevalueer vir kwaliteit, veiligheid of voorgestelde gebruik nie.

PPAT020/01